


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07D 495/04, 471/04, A61K 31/445</p>	<p>A1</p>	<p>(11) International Publication Number: <b>WO 99/43681</b> (43) International Publication Date: 2 September 1999 (02.09.99)</p>
<p>(21) International Application Number: PCT/US99/04185 (22) International Filing Date: 26 February 1999 (26.02.99) (30) Priority Data: 60/076,006 26 February 1998 (26.02.98) US (71) Applicant (for all designated States except US): NEUROGEN CORPORATION (US/US); 35 Northeast Industrial Road, Branford, CT 06405 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CAI, Guolin [CN/US]; 38 Gray Ledge Drive, Guilford, CT 06437 (US). LIU, Gang [CN/US]; 491 Pesaro Street, Agoura, CA 91301 (US). (74) Agent: SARUSSI, Steven, J.; McDonnell Boehnen Hulbert &amp; Berghoff, Suite 3200, 300 South Wacker Drive, Chicago, IL 60606 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: 4-(4-PIPERIDYLMETHYLAMINO) SUBSTITUTED HETEROARYL FUSED PYRIDINES: GABA BRAIN RECEPTOR LIGANDS</p> <p>(57) Abstract</p> <p>Disclosed are compounds of a formula (a) or the pharmaceutically acceptable non-toxic salts thereof wherein: R is hydrogen, alkyl, or (un)substituted alkoxy or amino; and W is (un)substituted alkyl, aryl, or heteroaryl, which compounds are highly selective agonists, antagonists or inverse agonists for GABA<sub>A</sub> brain receptors or prodrugs of agonists, antagonists or inverse agonists for GABA<sub>A</sub> brain receptors. The compounds of the invention are useful in the diagnosis and treatment of anxiety, Down Syndrome, sleep, cognitive and seizure disorders, and overdose with benzodiazepine drugs and for enhancement of alertness.</p> <div style="text-align: right;">  <p>(a)</p> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**4-(4-Piperidylmethylamino) Substituted Heteroaryl Fused  
Pyridines: GABA Brain Receptor Ligands**

5

**BACKGROUND OF THE INVENTION**

**Field of the Invention**

This invention relates to certain heterocyclic amino substituted heteroaryl fused pyridines which selectively bind to GABA<sub>A</sub> receptors. This invention also relates to pharmaceutical compositions comprising such compounds. It further relates to the use of such compounds in treating anxiety, sleep and seizure disorders, and overdoses of benzodiazepine-type drugs, and enhancing alertness. The interaction of heterocyclic amino substituted heteroaryl fused pyridines of the invention with a GABA binding site, the benzodiazepines (BDZ) receptor, is described. This interaction results in the pharmacological activities of these compounds.

20

**Description of the Related Art**

γ-Aminobutyric acid (GABA) is regarded as one of the major inhibitory amino acid transmitters in the mammalian brain. Over 40 years have elapsed since its presence in the brain was demonstrated (Roberts & Frankel, J. Biol. Chem. 187: 55-63, 1950; Udenfriend, J. Biol. Chem. 187: 65-69, 1950). Since that time, an enormous amount of effort has been devoted to implicating GABA in the etiology of seizure disorders, sleep, anxiety and cognition (Tallman and Gallager, Ann. Rev. Neuroscience 8: 21-44, 1985). Widely, although unequally, distributed through the mammalian brain, GABA is said to be a transmitter at approximately 30% of the synapses in the brain. GABA mediates many of its actions through a complex of proteins localized both on cell bodies and nerve endings; these are called GABA<sub>A</sub> receptors. Postsynaptic responses to

GABA are mediated through alterations in chloride conductance that generally, although not invariably, lead to hyperpolarization of the cell. Drugs that interact at the GABA<sub>A</sub> receptor can possess a spectrum of pharmacological activities depending on their abilities to modify the actions of GABA.

The 1,4-Benzodiazepines, such as diazepam, continue to be among the most widely used drugs in the world as anxiolytics, sedative-hypnotics, muscle relaxants, and anticonvulsants. A number of these compounds are extremely potent drugs; such potency indicates a site of action with a high affinity and specificity for individual receptors. Early electrophysiological studies indicated that a major action of benzodiazepines was enhancement of GABAergic inhibition. Presently, those compounds possessing activity similar to the benzodiazepines are called agonists. Compounds possessing activity opposite to benzodiazepines are called inverse agonists, and the compounds blocking both types of activity have been termed antagonists.

The GABA<sub>A</sub> receptor subunits have been cloned from bovine and human cDNA libraries (Schoenfield et al., 1988; Duman et al., 1989). A number of distinct cDNAs were identified as subunits of the GABA<sub>A</sub> receptor complex by cloning and expression. These are categorized into  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ ,  $\epsilon$ , and provide a molecular basis for the GABA<sub>A</sub> receptor heterogeneity and distinctive regional pharmacology (Shivvers et al., 1980; Levitan et al., 1989). The  $\gamma$  subunit appears to enable drugs like benzodiazepines to modify the GABA responses (Pritchett et al., 1989). The presence of low Hill coefficients in the binding of ligands to the GABA<sub>A</sub> receptor indicates unique profiles of subtype specific pharmacological action.


With the discovery of the "receptor" for the benzodiazepines and the subsequent definition of the nature of

the interaction between GABA and the benzodiazepines, it appears that the behaviorally important interactions of the benzodiazepines with different neurotransmitter systems are due in a large part to the enhanced ability of GABA itself to  
5 modify these systems. Each modified system, in turn, may be associated with the expression of a behavior. Depending on the mode of interaction, these compounds are capable of producing a spectrum of activities (either sedative, anxiolytic, and anticonvulsant, or wakefulness, seizures, and anxiety).

SUMMARY OF THE INVENTION

This invention provides novel compounds of Formula I which interact with a GABA<sub>A</sub> binding site, the benzodiazepine receptor.

- 5 The invention provides pharmaceutical compositions comprising compounds of Formula I. The invention also provides compounds useful in the diagnosis and treatment of anxiety, sleep and seizure disorders, overdose with benzodiazepine drugs and for enhancement of memory.
- 10 Accordingly, a broad embodiment of the invention is directed to compounds of general Formula I:



wherein:

- 15 the C ring represents a thiophene, pyridine, pyrazine, pyridazine, or pyrimidine ring, each of which is optionally substituted independently with one or two groups selected from C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, amino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkyl amino, or
- 20 trifluoromethyl;
- W is aryl, heteroaryl, 2- or 3-thienyl, or 2-, 3-, or 4-pyridyl, each of which may be mono, di- or trisubstituted independently with halogen, hydroxyl, lower alkyl, lower alkoxy, amino, or mono- or dialkylamino where each alkyl
- 25 portion is lower alkyl; or CONR<sub>4</sub>R<sub>5</sub> wherein R<sub>4</sub> and R<sub>5</sub> are the same or different and represent lower alkyl; and
- R represents hydrogen, lower alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, or aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl; or


R represents  $\text{CONR}_1\text{R}_2$  where  $\text{R}_1$  and  $\text{R}_2$  are the same or different and represent, hydrogen or lower alkyl or NR,  $\text{R}_2$  forms a ring having from 3-7, preferably 5-6, members.

5        These compounds are highly selective agonists, antagonists or inverse agonists for GABA<sub>A</sub> brain receptors or prodrugs of agonists, antagonists or inverse agonists for GABA<sub>A</sub> brain receptors. In other words, while the compounds of the invention all interact with GABA<sub>A</sub> brain receptors, they do  
10 not display identical physiological activity. Thus, these compounds are useful in the diagnosis and treatment of anxiety, sleep and seizure disorders, overdose with benzodiazepine drugs and for enhancement of memory. For example, these compounds can be used to treat overdoses of  
15 benzodiazepine-type drugs as they would competitively bind to the benzodiazepine receptor.

DETAILED DESCRIPTION OF THE INVENTION

The novel compounds encompassed by the instant invention can be described by general Formula I set forth above or the pharmaceutically acceptable non-toxic salts thereof.

5 In addition, the present invention also encompasses compounds of Formula IIa and IIb



or the pharmaceutically acceptable non-toxic salts thereof  
 10 wherein W and R are as defined above for Formula I; and R<sub>a</sub> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> alkoxy, halogen, hydroxy, amino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkyl amino, or trifluoromethyl.


Preferred compounds of Formula IIa and IIb are where W is  
 15 aryl or heteroaryl mono or disubstituted independently with halogen, hydroxyl, lower alkyl, or lower alkoxy.

Other preferred compounds of Formula IIa and IIb are where R is hydrogen, lower alkyl.

Preferred R<sub>a</sub> groups include C<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy,  
 20 hydroxy, dimethyl amino, diethylamino, ethylamino, chloro, fluoro and bromo. Particularly preferred R<sub>a</sub> groups are chloro, bromo, fluoro, methylethyl and amino.

The present invention also encompasses compounds of Formula III:






III

wherein W and R are as defined above for Formula I; and  
 A, B, C, and D are independently CR<sub>3</sub>, or nitrogen, provided  
 5 that no more than two of A, B, C, and D are nitrogen  
 simultaneously;  
 R<sub>3</sub> is hydrogen, alkyl, halogen, hydroxy, or lower alkoxy.

Other preferred compounds of the invention are  
 10 encompassed by the following formulae:




IV

where

R is as defined above for Formula I; and  
 15 W is aryl or 2-, 3-, or 4-pyridyl, each of which may be  
 mono or disubstituted independently with halogen,  
 lower alkyl, lower alkoxy, amino, or mono- or  
 dialkylamino where each alky portion is lower alkyl,  
 or

CONR<sub>4</sub>R<sub>5</sub> wherein R<sub>4</sub> and R<sub>5</sub> are same or different and represent lower alkyl.

More preferred compounds of Formula IV are those where W  
5 is



wherein

R<sub>6</sub> is halogen, hydroxy, lower alkyl, lower alkoxy, amino, or mono- or dialkylamino where each alkyl portion is  
10 lower alkyl, or


R<sub>6</sub> is CONR<sub>4</sub>R<sub>5</sub> wherein R<sub>4</sub> and R<sub>5</sub> are the same or different and represent lower alkyl; and

X is CR<sub>2</sub> or nitrogen, where R<sub>2</sub> is hydrogen, lower alkyl, halogen, hydroxy, or lower alkoxy.

15

Even more preferred compounds of Formula IV are those where R is hydrogen, R<sub>6</sub> is hydrogen or lower alkoxy, and X is CR<sub>2</sub> or nitrogen, where R<sub>2</sub> is hydrogen, fluorine, methoxy, or ethoxy.

20 The present invention also encompasses compounds of Formula V:




where

25 R is as defined above for Formula I; and

W is aryl or 2-, 3-, or 4-pyridyl, each of which may be  
 mono or disubstituted independently with halogen,  
 lower alkyl, lower alkoxy, amino, or mono- or  
 dialkylamino where each alkyl portion is lower  
 5 alkyl, or  $\text{CONR}_4\text{R}_5$  where  $\text{R}_4$  and  $\text{R}_5$  are same or  
 different and represent lower alkyl.

More preferred compounds of Formula V are those where W  
 is




10

wherein

$\text{R}_6$  is halogen, hydroxy, lower alkyl, lower alkoxy, amino,  
 or mono- or dialkylamino where each alkyl portion is  
 lower alkyl, or  $\text{CONR}_7\text{R}_8$  wherein  $\text{R}_7$  and  $\text{R}_8$  are the  
 15 same or different and represent lower alkyl; and  
 $\text{X}$  is  $\text{CR}_9$  or nitrogen, where  $\text{R}_9$  is hydrogen, lower alkyl,  
 halogen, hydroxy, or lower alkoxy.

Even more preferred compounds of Formula V are those  
 20 where  $\text{R}$  is hydrogen,  $\text{R}_6$  is hydrogen or lower alkoxy, and  $\text{X}$  is  
 $\text{CR}_9$  or nitrogen, where  $\text{R}_9$  is hydrogen, fluorine, methoxy, or  
 ethoxy.

The present invention also encompassess compounds of  
 Formula VI:



25

## VI

where


A, B, C, and D are CR<sub>1</sub>, or nitrogen, provided that no more than two of A, B, C, and D are nitrogen simultaneously;

R<sub>1</sub> is hydrogen, lower alkyl, halogen, hydroxy, or lower alkoxy;

R is as defined above for Formula I; and

W is aryl or 2-, 3-, or 4-pyridyl, each of which may be mono or disubstituted independently with halogen, hydroxy, lower alkyl, lower alkoxy, amino, or mono- or dialkylamino where each alkyl portion is lower alkyl, or CONR<sub>4</sub>R<sub>5</sub> wherein R<sub>4</sub> and R<sub>5</sub> are same or different and representing lower alkyl.

More preferred compounds of Formula VI are those where W is



wherein

R<sub>6</sub> is halogen, hydroxy, lower alkyl, lower alkoxy, amino, mono- or dialkylamino where each alkyl portion is lower alkyl, or CONR<sub>7</sub>R<sub>8</sub> wherein R<sub>7</sub> and R<sub>8</sub> are the same or different and represent lower alkyl; and

X is CR<sub>1</sub>, or nitrogen, and R<sub>1</sub> is hydrogen, lower alkyl, halogen, hydroxy, or lower alkoxy.

Even more preferred compounds of Formula VI are those where A is nitrogen, B, C, and D are hydrogen, R is hydrogen, R<sub>2</sub> is hydrogen, and X is CR<sub>1</sub>, where R<sub>1</sub> is hydrogen, fluorine, methoxy or ethoxy.

Representative compounds of the present invention, which are encompassed by Formula I, include, but are not limited to

the compounds in Table I and their pharmaceutically acceptable acid and base addition salts. In addition, if the compound of the invention is obtained as an acid addition salt, the free base can be obtained by basifying a solution of the acid salt.

5   Conversely, if the product is a free base, an addition salt, particularly a pharmaceutically acceptable addition salt, may be produced by dissolving the free base in a suitable organic solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts  
10   from base compounds.

Non-toxic pharmaceutically acceptable salts include salts of acids such as hydrochloric, phosphoric, hydrobromic, sulfuric, sulfinic, formic, toluenesulfonic, methanesulfonic, nitric, benzoic, citric, tartaric, maleic, hydroiodic,  
15   alkanoic such as acetic,  $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$  where  $n$  is 0-4, and the like. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.

The present invention also encompasses the acylated  
20   prodrugs of the compounds of Formula I. Those skilled in the art will recognize various synthetic methodologies which may be employed to prepare non-toxic pharmaceutically acceptable addition salts and acylated prodrugs of the compounds encompassed by Formula I.

25   By "alkyl" and "lower alkyl" in the present invention is meant straight or branched chain alkyl groups having 1-6 carbon atoms, such as, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, and 3-methylpentyl.

30   By "alkoxy" and "lower alkoxy" in the present invention is meant straight or branched chain alkoxy groups having 1-6 carbon atoms, such as, for example, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-

butoxy, pentoxy, 2-pentyl, isopentoxy, neopentoxy, hexoxy, 2-hexoxy, 3-hexoxy, and 3-methylpentoxy.



By the term "halogen" in the present invention is meant fluorine, bromine, chlorine, and iodine.


5 By heteroaryl is meant one or more aromatic ring systems of 5-, 6-, or 7-membered rings containing at least one and up to four heteroatoms selected from nitrogen, oxygen, or sulfur. Such heteroaryl groups include, for example, thienyl, furanyl, thiazolyl, imidazolyl, (is)oxazolyl, pyridyl, pyrimidinyl,  
10 (iso)quinolinyl, naphthyridinyl, benzimidazolyl, benzoxazolyl.

By aryl is meant an aromatic carbocyclic group having a single ring (e.g., phenyl), multiple rings (e.g., biphenyl), or multiple condensed rings in which at least one is aromatic, (e.g., 1,2,3,4-tetrahydronaphthyl, naphthyl, anthryl, or  
15 phenanthryl), which is optionally mono-, di-, or trisubstituted with, e.g., halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, lower acyloxy, aryl, heteroaryl, and hydroxy.


Representative compounds of the invention are shown below  
20 in Table 1.

Table 1







Compound 4




Compound 5




Compound 6



Compound 7



Compound 8



Compound 9

The pharmaceutical utility of compounds of this invention is indicated by the assays for GABA<sub>A</sub> receptor binding activity set forth in the Examples below.

The compounds of Formula I and their salts are suitable for the diagnosis and treatment of anxiety, Down Syndrome, sleep, cognitive and seizure disorders, and overdose with benzodiazepine drugs and for enhancement of alertness, both in  
5 human and non-human animals and domestic pets, especially dogs and cats and farm animals such as sheep, swine and cattle.

The compounds of general Formula I may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional  
10 non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a  
15 compound of general Formula I and a pharmaceutically acceptable carrier. One or more compounds of general Formula I may be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The  
20 pharmaceutical compositions containing compounds of general Formula I may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

25 Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and  
30 preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture



of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or  
5 alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract  
10 and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with  
15 an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in  
20 admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia;  
25 dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example  
30 heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters

derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one  
5 or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil  
10 such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the  
15 addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives.  
20 Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

Pharmaceutical compositions of the invention may also be  
25 in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring  
30 phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for

example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or  
5 sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those  
10 suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among  
15 the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono-  
20 or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of general Formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be  
25 prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

30 Compounds of general Formula I may be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as

local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.


Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 1 mg to about 500 mg of an active ingredient.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

For administration to non-human animals, the composition may also be added to the animal feed or drinking water. It will be convenient to formulate these animal feed and drinking water compositions with a mullet-dose of the drug so that the animal takes in an appropriate quantity of the composition along with its diet. It will also be convenient to present the composition as a premix for addition to the feed or drinking water.

An illustration of the preparation of compounds of the present invention is given in Schemes I.

Scheme I



In the above scheme, the W and C rings are as defined  
 5 above for Formula I.

An illustration of the preparation of compounds of the  
 present invention is given in Scheme I. Those having skill in  
 the art will recognize that the starting materials may be  
 varied and additional steps employed to produce compounds  
 10 encompassed by the present invention, as demonstrated by the  
 following examples.

As shown, an aniline is reacted with a  $\beta$ -keto ester  
 in the presence of an acid, such as, for example, P-  
 toluenesulfonic acid, to form a 4-hydroxypyridine which is  
 15 subsequently converted to the 4-chloropyridine upon treatment  
 with a nucleophilic halogenating reagent such as phosphorus  
 oxychloride. The resulting chloride is reacted with 4-  
 (Aminomethyl)piperidine at elevated temperatures to form the N-  
 alkylated product. The piperidine can then be further  
 20 derivatized with a desired isocyanate, such as, for example,  
 methyl isocyanate, to form the target compound.

The disclosures in this application of all articles and references, including patents, are incorporated herein by reference.

The invention is illustrated further by the following  
5 examples which are not to be construed as limiting the invention in scope or spirit to the specific procedures described in them.


The starting materials and various intermediates may be obtained from commercial sources, prepared from commercially  
10 available organic compounds, or prepared using well known synthetic methods.

Representative examples of methods for preparing intermediates of the invention are set forth below.


15

EXAMPLE 1

1. 5-(4-Fluorophenyl)-thieno[3,2-b]pyridin-7-ol




A mixture of 3-amino-2-thiophenecarboxylic acid (8 g, 49 mmol), ethyl 4-fluorobenzoylacetate (9.6 g, 49 mmol), and p-toluenesulfonic acid monohydrate (0.2 g, 1 mmol) in toluene  
20 (100 mL) is refluxed for 20 hours with a Dean-Stark water trap to remove produced water. After cooling to room temperature, the precipitate is filtered and washed with diethyl ether. The solid is dissolved in diphenyl ether (80 mL) and heated at  
25 220°C for 2 hours. The reaction solution is then cooled to room temperature, diethyl ether is added, and the precipitate is filtered and washed with diethyl ether to give 5-(4-fluorophenyl)-thieno[3,2-b]pyridin-7-ol (2 g, 17% yield) as brown crystalline needles, m.p. 316-318 °C.

2. 7-Chloro-5-(4-fluorophenyl)thieno[3,2-b]pyridine

A solution of 5-(4-fluorophenyl)-thieno[3,2-b]pyridin-7-ol (1.6 g) in phosphorus oxychloride (50 mL) is refluxed for 3 hours. After the excess amount of phosphorus oxychloride is removed under vacuum, the residue is treated with ethyl acetate (20 mL), and NaOH (2N, 20 mL). The mixture is extracted with ethyl acetate (3x20 mL). The combined organic layers are washed with brine and dried over MgSO<sub>4</sub>. Evaporation of the solvent affords 7-chloro-5-(4-fluorophenyl)thieno[3,2-b]pyridine (1.5 g, 88% yield) as a white solid, m.p. 119-121 °C.

3. N-(4-piperidinylmethyl)-5-(4-fluorophenyl)thieno[3,2-b]pyridin-7-amine

A solution of 7-chloro-5-(4-fluorophenyl)thieno[3,2-b]pyridine (60 mg, 0.23 mmole) in 4-aminomethylpiperidine (1 mL) is heated at 160 °C oil bath under N<sub>2</sub> for 4 hours. The reaction mixture is cooled to room temperature and purified on preparative tlc plate to give N-(4-piperidinylmethyl)-5-(4-fluorophenyl)thieno[3,2-b]pyridin-7-amine (15 mg, 19% yield) as a colorless oil. This material is dissolved in 1 mL ethyl acetate. Ethyl acetate saturated with HCl (2 mL) is added and then concentrated to afford N-(4-piperidinylmethyl)-5-(4-fluorophenyl)thieno[3,2-b]pyridin-7-amine dihydrochloride (compound 1) as a greasy oil. The salt is triturated with ether, collected by filtration, washed with ether and dried in vacuo, m.p. >260°C (dec).

**EXAMPLE 2**4-Chloro-6-phenylthieno[2,3-b]pyridine


5 A mixture of 3-aminothiophene (4 g, 0.04 mole), ethyl benzoylacetate (11 mL, 0.06 mole), toluene (100 mL), and p-toluene sulfonic acid monohydrate (300 mg) is stirred and heated under reflux with a Dean-Stark water trap at 120 °C for 16 hours. The reaction mixture is concentrated, and diphenyl  
10 ether (30 mL) is added. After heating at 240 °C for 1 hour, the reaction mixture was allowed to cool to room temperature. The reaction mixture is diluted with hexane, and the semi-solid collected is then purified by short silica gel column (CH<sub>2</sub>Cl<sub>2</sub> / MeOH / NH<sub>4</sub>OH: 10:1:0.1). The resulting product, 4-  
15 hydroxy-6-phenyl-thieno[2,3-b]pyridine, is treated with phosphorus oxychloride (30 mL) and heated under reflux for 3 hours. This mixture is cooled to room temperature, poured onto ice, neutralized with 10 N sodium hydroxide, and extracted with methylene chloride (3x20 mL). The combined  
20 organic layers are dried over sodium sulfate, concentrated in vacuo. and the residue is purified by chromatography to give 4-chloro-6-phenylthieno[2,3-b]pyridine as a yellowish solid (1.1 g, 11% total yield), m.p. 83-85 °C.

25

**EXAMPLE 3**

N-(4-N-ethylcarboxamidopiperidinylmethyl)-5-(4-fluorophenyl)thieno[3,2-b]pyridin-7-amine





Compound 2

A reaction solution of N-(4-piperidinylmethyl)-5-(4-fluorophenyl)thieno[3,2-b]pyridin-7-amine (10 mg, 0.03 mmole), ethyl isocyanate (0.1 mL) in toluene (5 mL) is heated at 120 °C oil bath for 1 hour. The resulting solution is cooled down to room temperature and concentrated. The residue is purified on a preparative tlc plate to give N-(4-N-ethylcarboxamidopiperidinylmethyl)-5-(4-fluorophenyl)thieno[3,2-b]pyridin-7-amine (1 mg, 8% yield) as a colorless oil. This material is dissolved in 1 mL ethyl acetate. Ethyl acetate saturated with HCl (2 mL) is added and then concentrated to afford N-(4-N-ethylcarboxamidopiperidinylmethyl)-5-(4-fluorophenyl)thieno[3,2-b]pyridin-7-amine hydrochloride (compound 2) as a greasy oil. The salt was solidified with ether.

#### EXAMPLE 4

The following compounds are prepared essentially according to the procedure described in Examples 1-5:

- a) N-(4-Piperidinylmethyl)-5-phenylthieno[3,2-b]pyridin-7-amine dihydrochloride (compound 3).

b) N-(4-Piperidinylmethyl)-5-(3-methoxyphenyl)thieno[3,2-b]pyridin-7-amine dihydrochloride (compound 4).

5 c) N-(4-Piperidinylmethyl)-6-phenylthieno[2,3-b]pyridin-4-amine dihydrochloride (compound 5).

d) N-(4-Piperidinylmethyl)-5-(4-pyridinyl)thieno[3,2-b]pyridin-7-amine dihydrochloride (compound 6).

10

e) N-(4-Piperidinylmethyl)-5-(4-ethoxyphenyl)thieno[3,2-b]pyridin-7-amine dihydrochloride (compound 7).

15 f) N-(4-N-Ethylcarboxamidopiperidinylmethyl)-5-(4-fluorophenyl)thieno[3,2-b]pyridin-7-amine hydrochloride (compound 8).

g) 4-(4-Piperidinylmethylamino)-2-phenyl-1,5-naphthyridine. (compound 9).

20

#### EXAMPLE 5

The pharmaceutical utility of compounds of this invention is indicated by the following assay for GABA<sub>A</sub> receptor binding activity.

25

Assays are carried out as described in Thomas and Tallman (J. Bio. Chem. 156: 9838-9842 , J. Neurosci. 3: 433-440, 1983). Rat cortical tissue is dissected and homogenized in 25 volumes (w/v) of 0.05 M Tris HCl buffer (pH 7.4 at 4°C). The tissue homogenate is centrifuged in the cold (4°C) at 20,000 x g for 20'. The supernatant is decanted and the pellet is rehomogenized in the same volume of buffer and again centrifuged at 20,000 x g. The supernatant is decanted and

30

the pellet is frozen at  $-20^{\circ}\text{C}$  overnight. The pellet is then thawed and rehomogenized in 25 volume (original wt/vol) of buffer and the procedure is carried out twice. The pellet is finally resuspended in 50 volumes (w/vol of 0.05M Tris HCl  
5 buffer (pH 7.4 at  $40^{\circ}\text{C}$ ).

Incubations contain 100ml of tissue homogenate, 100ml of radioligand 0.5nM ( $^3\text{H}$ -RO15-1788 [ $^3\text{H}$ -Flumazenil] specific activity 80 Ci/mmol), drug or blocker and buffer to a total volume of 500 ml. Incubations are carried for 30min at  $4^{\circ}\text{C}$   
10 then are rapidly filtered through GFB filters to separate free and bound ligand. Filters are washed twice with fresh 0.05M Tris HCl buffer (pH 7.4 at  $4^{\circ}\text{C}$ ) and counted in a liquid scintillation counter. 1.0mM diazepam is added to some tubes to determine nonspecific binding. Data are collected in  
15 triplicate determinations, averaged and % inhibition of total specific binding is calculated. Total Specific Binding = Total - Nonspecific. In some cases, the amounts of unlabeled drugs is varied and total displacement curves of binding are carried out. Data are converted to  $\text{IC}_{50}$  or  $\text{K}_i$ . The  $\text{K}_i$  values  
20 for the compounds in this invention are less than 200 nM.

#### EXAMPLE 6

In addition, the following assay may be used to determine if the compounds of the invention are agonists, antagonists,  
25 or inverse agonists, and, therefore, their specific pharmaceutical utility. The following assay can be employed to determine specific GABA<sub>A</sub> receptor activity.

Assays are carried out as described in White and Gurley (NeuroReport 6: 1313-1316, 1995) and White, Gurley, Hartnett,  
30 Stirling, and Gregory (Receptors and Channels 3: 1-5, 1995) with modifications. *Xenopus Laevis* oocytes are enzymatically isolated and injected with non-polyadenylated cRNA mixed in a

ratio of 4:1:4 for human derived  $\alpha$ ,  $\beta$ , and  $\gamma$  subunits, respectively. For each subunit combination, sufficient message is injected to result in current amplitudes of  $>10$  nA when  $1 \mu\text{M}$  GABA is applied.

5 Electrophysiological recordings are carried out using the two electrode voltage-clamp technique at a membrane holding potential of  $-70$  mV.

Compounds are evaluated against a GABA concentration that evokes  $<10\%$  of the maximal evokable GABA current. Each oocyte  
10 is exposed to increasing concentrations of compound in order to evaluate a concentration/effect relationship. Compound efficacy is expressed as a percent-change in current amplitude:  $100 \cdot ((I_c/I) - 1)$ , where  $I_c$  is the GABA evoked current amplitude observed in the presence of compound and  $I$  is the  
15 GABA evoked current amplitude observed in the absence of compound.

Specificity of a compound for the Ro15-1788 site is determined following completion of the concentration/effect curve. After washing the oocyte sufficiently to remove  
20 previously applied compound, the oocyte is exposed to GABA +  $1 \mu\text{M}$  Ro15-1788, followed by exposure to GABA +  $1 \mu\text{M}$  Ro15-1788 + compound. Percent change due to addition of compound is calculated as described above. Any percent change observed in the presence of Ro15-1788 is subtracted from the percent  
25 changes in current amplitude observed in the absence of  $1 \mu\text{M}$  Ro15-1788. These net values are used for the calculation of average efficacy and  $\text{EC}_{50}$  values.


To evaluate average efficacy and  $\text{EC}_{50}$  values, the concentration/effect data are averaged across cells and fit to  
30 the logistic equation. Average values are reported as mean  $\pm$  standard error.

The invention and the manner and process of making and using it, are now described in such full, clear, concise and

exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to be understood that the foregoing describes preferred embodiments of the present invention and that modifications may be made  
5 therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as invention, the following claims conclude this specification.

## WHAT IS CLAIMED IS:

1. A compound of the formula:



or the pharmaceutically acceptable non-toxic salts thereof  
5 wherein:

the C ring represents a thiophene, pyridine, pyrazine,  
pyridazine, or pyrimidine ring each of which is  
optionally substituted with one or two groups  
independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy,  
10 halogen, hydroxy, amino, mono- or di-(C<sub>1</sub>-C<sub>6</sub>)alkylamino, or  
trifluoromethyl;


W is aryl, heteroaryl, 2- or 3-thienyl, or 2-, 3-, or 4-  
pyridyl, each of which is optionally mono or  
disubstituted independently with halogen, hydroxyl, lower  
15 alkyl, lower alkoxy, amino, or mono- or dialkylamino  
where each alkyl portion is lower alkyl; or CONR<sub>4</sub>R<sub>5</sub>  
wherein R<sub>4</sub> and R<sub>5</sub> are the same or different and represent  
lower alkyl; and

R represents hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
20 amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-  
C<sub>6</sub>)alkyl, alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryl(C<sub>1</sub>-  
C<sub>6</sub>)alkyl; or

CONR<sub>1</sub>R<sub>2</sub> wherein R<sub>1</sub> and R<sub>2</sub> are the same or different and  
represent hydrogen or lower alkyl.

25

2. A compound according to Claim 1 which is:




or the pharmaceutically acceptable non-toxic salts thereof wherein:

$R_1$  is hydrogen,  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  alkoxy, halogen, hydroxy, amino, mono- or di( $C_1$ - $C_6$ )alkyl amino, or trifluoromethyl;

$R$  is hydrogen or lower alkyl; and

$W$  is aryl, heteroaryl, 2- or 3-thienyl, or 2-, 3-, or 4-pyridyl, each of which may be mono or disubstituted independently with halogen, hydroxyl, lower alkyl, lower alkoxy, amino, or mono- or dialkylamino where each alkyl portion is lower alkyl or  $CONR_4R_5$  wherein  $R_4$  and  $R_5$  are the same or different and represent lower alkyl.

3. A compound according to Claim 1 which is:




or the pharmaceutically acceptable non-toxic salts thereof wherein:

$R_a$  is hydrogen,  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  alkoxy, halogen, hydroxy, amino, mono- or di( $C_1$ - $C_6$ )alkyl amino, or trifluoromethyl

R is hydrogen or lower alkyl; and

W is aryl, heteroaryl, 2- or 3-thienyl, or 2-, 3-, or 4-pyridyl, each of which may be mono or disubstituted independently with halogen, hydroxyl, lower alkyl, lower alkoxy, amino, or mono- or dialkylamino where each alkyl portion is lower alkyl.

4. A compound according to Claim 1 which is:



10


or the pharmaceutically acceptable non-toxic salts thereof wherein:

W is aryl, heteroaryl, 2- or 3-thienyl, or 2-, 3-, or 4-pyridyl, each of which may be mono or disubstituted independently with halogen, hydroxyl, lower alkyl, lower alkoxy, amino, or mono- or dialkylamino where each alkyl portion is lower alkyl; or  $CONR_4R_5$ , wherein  $R_4$  and  $R_5$  are the same or different and represent lower alkyl and

$R_2$  represents  
hydrogen, lower alkyl having, hydroxyalkyl, aminoalkyl, alkoxyalkyl, lower alkoxy, or arylalkyl where each alkyl portion is lower alkyl.

5. A compound of the formula:





or the pharmaceutically acceptable non-toxic salts thereof wherein:


5 R is hydrogen, lower alkyl, hydroxyalkyl, aminoalkyl, alkoxyalkyl, lower alkoxy, or arylalkyl where each alkyl portion is lower alkyl; or  $\text{CONR}_1\text{R}_2$  wherein  $\text{R}_1$  and  $\text{R}_2$  are the same or different and represent lower alkyl;

10  $\text{R}_6$  is halogen, hydroxy, lower alkyl, lower alkoxy, amino, mono- or dialkylamino where each alkyl portion is lower alkyl, or  $\text{CONR}_4\text{R}_5$  wherein  $\text{R}_4$  and  $\text{R}_5$  are the same or different and represent lower alkyl; and

X is nitrogen, or  $\text{CR}$ , wherein  $\text{R}$  is hydrogen, alkyl, halogen, hydroxy, or lower alkoxy.

15

6. A compound of the formula:




or the pharmaceutically acceptable non-toxic salts thereof wherein:

5 R is hydrogen, lower alkyl, hydroxyalkyl, aminoalkyl, alkoxyalkyl, lower alkoxy, arylalkyl where each alkyl portion is lower alkyl, or  $\text{CONR}_1\text{R}_2$  wherein  $\text{R}_1$  and  $\text{R}_2$  are the same or different and represent lower alkyl;

10  $\text{R}_6$  is halogen, hydroxy, lower alkyl, lower alkoxy, amino, mono- or dialkylamino where each alkyl portion is lower alkyl, or  $\text{CONR}_4\text{R}_5$  wherein  $\text{R}_4$  and  $\text{R}_5$  are the same or different and represents lower alkyl; and

X is nitrogen, or  $\text{CR}$ , wherein  $\text{R}$  is hydrogen, alkyl, halogen, hydroxy, or lower alkoxy.

15 7. A compound of the formula:



or the pharmaceutically acceptable non-toxic salts thereof wherein:

20 A, B, C, and D are  $\text{CR}$ , or nitrogen, provided that no more than two of A, B, C, or D are nitrogen simultaneously;

$\text{R}_6$  is hydrogen, alkyl, halogen, hydroxy, or lower alkoxy having;

25 R is hydrogen, lower alkyl, hydroxyalkyl, aminoalkyl, alkoxyalkyl, lower alkoxy, arylalkyl where each

alkyl portion is lower alkyl, or  $\text{CONR}_1\text{R}_2$  wherein  $\text{R}_1$  and  $\text{R}_2$  are the same or different and represent lower alkyl;

$\text{R}_6$  is halogen, hydroxy, lower alkyl, lower alkoxy, amino, or mono- or dialkylamino where each alkyl portion is lower alkyl, or  $\text{CONR}_4\text{R}_5$  wherein  $\text{R}_4$  and  $\text{R}_5$  are the same or different and represent lower alkyl; and

$\text{X}$  is  $\text{CR}_7$  or nitrogen, where  $\text{R}_7$  is hydrogen, lower alkyl, halogen, hydroxy, or lower alkoxy.

10

8. A compound according to Claim 1 which is N-(4-piperidinylmethyl)-5-(4-fluorophenyl)thieno[3,2-b]pyridin-7-amine.

15

9. A compound according to Claim 1 which is N-(4-piperidinylmethyl)-5-phenylthieno[3,2-b]pyridin-7-amine.

20

10. A compound according to Claim 1 which is N-(4-piperidinylmethyl)-5-(3-methoxyphenyl)thieno[3,2-b]pyridin-7-amine.

25

11. A compound according to Claim 1 which is N-(4-piperidinylmethyl)-5-(4-ethoxyphenyl)thieno[3,2-b]pyridin-7-amine.

12. A compound according to Claim 1 which is N-(4-piperidinylmethyl)-5-(4-pyridinyl)thieno[3,2-b]pyridin-7-amine.

30

13. A compound according to Claim 1 which is N-(4-piperidinylmethyl-N-ethylcarboxamide)-5-(4-fluorophenyl)thieno[3,2-b]pyridin-7-amine.

14. A compound according to Claim 1 which is 8-(N-(piperidin-4-ylmethyl)amino)-2-phenyl-1,5-naphthyridine.

5 15. A compound according to Claim 1 which is N-(4-piperidinylmethyl)-6-phenylthieno[2,3-b]pyridin-4-amine.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/04185

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D495/04 C07D471/04 A61K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 02433 A (NEUROGEN CORPORATION) 22 January 1998 (1998-01-22) the whole document ---	1
A	WO 98 02420 A (NEUROGEN CORPORATION) 22 January 1998 (1998-01-22) the whole document ---	1
A	WO 95 11885 A (NEUROGEN CORPORATION) 4 May 1995 (1995-05-04) the whole document ---	1
A	US 5 243 049 A (KENNETH SHAW ET AL.) 7 September 1993 (1993-09-07) the whole document --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

27 July 1999

Date of mailing of the international search report

03/08/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Kyriakakou, G

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/04185

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GB 1 177 849 A (STERLING DRUG INC.) 14 January 1970 (1970-01-14) the whole document ---	1
A	LUISA SAVINI ET AL.: "Pyrazolo'4,3-c! quinolines: Synthesis and specific inhibition of Benzodiazepine Receptor binding" IL FARMACO, vol. 48, no. 1, 1993, pages 65-75, XP002081964 the whole document ---	1
A	CHRISTOPHER D. BENHAM ET AL.: "The synthesis of pyrazolo'4,3-c!- and imidazo'4,5-c!-aryl'e!fused pyridines as structural analogues of 4-aminonicotinates" BIOORGANIC AND MEDICINAL CHEMISTRY LETTERS, vol. 5, no. 21, 1995, pages 2455-2460, XP004136805 great britain the whole document ---	1
A	T. P. BLACKBURN ET AL.: "Isosteric Replacement of the Indole nucleus by Benzothiophene in a series of Pyrido(2,3-b!indoles with potential anxiolytic activity" BIOORGANIC AND MEDICINAL CHEMISTRY LETTERS, vol. 5, no. 22, 1995, pages 2589-2592, XP004135197 great britain the whole document -----	1

## INTERNATIONAL SEARCH REPORT

Inter national Application No

PCT/US 99/04185

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9802433 A	22-01-1998	US 5750702 A	12-05-1998
		AU 2832897 A	09-02-1998
		AU 3660497 A	09-02-1998
		HR 970385 A	31-08-1998
		WO 9802420 A	22-01-1998
WO 9802420 A	22-01-1998	US 5750702 A	12-05-1998
		AU 2832897 A	09-02-1998
		AU 3660497 A	09-02-1998
		HR 970385 A	31-08-1998
		WO 9802433 A	22-01-1998
WO 9511885 A	04-05-1995	US 5484944 A	16-01-1996
		AT 168673 T	15-08-1998
		AU 701151 B	21-01-1999
		AU 8126594 A	22-05-1995
		CA 2175204 A	11-05-1995
		CN 1143953 A	26-02-1997
		CZ 9601129 A	16-04-1997
		DE 69411939 D	27-08-1998
		DE 69411939 T	25-03-1999
		EP 0725775 A	14-08-1996
		EP 0825193 A	25-02-1998
		ES 2122504 T	16-12-1998
		FI 961776 A	14-06-1996
		HU 76064 A	30-06-1997
		JP 9506082 T	17-06-1997
		NO 961655 A	10-06-1996
		NZ 275968 A	27-05-1998
		PL 314136 A	19-08-1996
		SE 9601563 A	22-05-1996
		SK 52496 A	09-07-1997
US 5243049 A	07-09-1993	US 5608079 A	04-03-1997
		US 5750702 A	12-05-1998
		US 5908932 A	01-06-1999
GB 1177849 A	14-01-1970	WO 9426741 A	24-11-1994
		US 5604235 A	18-02-1997
		BE 706646 A	16-05-1968
		FR 7325 M	06-10-1969
		NL 6715624 A	24-05-1968
		US 3505339 A	07-04-1970